## PATENT ABSTRACTS OF JAPAN

CO7C 91/15

(11)Publication number:

01-093562

(43) Date of publication of application: 12.04.1989

(51)Int.CI.

A61K 31/13 A61K 31/215 A61K 31/66 A61K 31/665 A61K 31/70 CO7C 93/22 C07C103/38 C07C103/50 C07C117/00 C07C125/065 C07C127/15 C07C127/19 C07C127/22 C07C143/75 C07C157/09 C07H 15/04 // C07C103/60 CO7F 7/18

(21)Application number: 62-252276

(22)Date of filing:

05.10.1987

(71)Applicant : SHIONOGI & CO LTD

(72)Inventor: SUGIMOTO HIROHIKO

SUGAWARA TAMIO MAKINO ITSUO SATO KOZABURO NARISADA MASAYUKI

## (54) PRODUCTION OF SPHINGOSINE DERIVATIVE

(57)Abstract:

NEW MATERIAL: The compound of formula I (R1 is H, acyl, glycosyl or phosphoric acid ester residue; R2 is H, acyl or glycosyl; A is (substituted) amino or azido; n is 10W14; excluding the compounds corresponding to known substances).

EXAMPLE: (2S, 3R)-2-acetamido-1,3-dihydroxy-4-cis-octadecene.

USE: Carcinostatic agent.

PREPARATION: The compound of formula I can be produced by acylating the amino group of the compound of formula II. The introduction of these groups is carried cut by reacting with corresponding carboxylic acid (e.g. acetic acid), acid anhydride (e.g. acetic anhydride) or acid chloride (e.g. acetyl chloride). For example, N-bydroxy-5-norbornene-2,3-dicarboximide is preferably used in the acylation by carbodiimide process and, in the case of acylation with acid anhydride, the reaction is preferably carried out in an alcoholic solvent.

## **LEGAL STATUS**

[Date of request for examination]

[Date of sending the examiner's decision of rejection]

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

[Date of registration]

[Number of appeal against examiner's decision of rejection]

[Date of requesting appeal against examiner's decision of rejection]

[Date of extinction of right]